



## Heart Failure and Cardiomyopathies

### PLASMA ENDOTHELIN-1 INDEPENDENTLY PREDICTS 180-DAY MORTALITY IN ACUTE HEART FAILURE: AN ASCEND-HF SUB-STUDY

Moderated Poster Contributions

Heart Failure and Cardiomyopathies Moderated Poster Theater, Poster Hall B1  
Saturday, March 14, 2015, 11:15 a.m.-11:25 a.m.

Session Title: Biomarkers in Heart Failure

Abstract Category: 14. Heart Failure and Cardiomyopathies: Clinical

Presentation Number: 1128M-13

Authors: Antonio Perez, Justin Grodin, Adrian Hernandez, Javed Butler, Marco Metra, G. Michael Felker, Adriaan A. Voors, John McMurray, Paul Armstrong, Robert Califf, Randall Starling, Christopher O'Connor, Wai Hong Tang, Cleveland Clinic, Cleveland, OH, USA

**Background:** Endothelin-1 (ET-1) is an endogenous vasoconstrictor implicated in pulmonary and systemic hypertension as well as ventricular dysfunction, through effects on vascular smooth muscle, neurohormonal vascular regulation and cardiomyocytes.

**Methods:** We measured baseline ET-1 level (Singulex) in acute decompensated heart failure (ADHF) and 180-day mortality, using a cohort of 872 patients from the ASCEND-HF trial (randomized to nesiritide vs placebo).

**Results:** Median ET-1 was 7.6 pg/mL (IQR 5.9-10) at baseline, 6.3 pg/mL (IQR 4.9-8.1) at 48-72 hours, and 5.9 pg/mL (IQR 4.7-7.9) at 30 days. At 180 days, 102 patients had died. Elevated baseline ET-1 was associated with increased 180-day mortality risk (Figure, log-transformed ET-1: HR 3.29, 95% CI 2.08-5.22,  $p < 0.0001$ ). Multivariable adjustments for age, log-transformed BUN, sodium, baseline hypotension and baseline dyspnea at rest also showed statistical significance (log-transformed ET-1: HR 2.74, 95% CI 1.68-4.45,  $p < 0.0001$ ). When log-transformed baseline NTproBNP was included as an additional covariate, ET-1 remained independently associated with 180-day mortality (log-transformed ET-1: HR 1.98, 95% CI 1.18-3.32,  $p = 0.009$ ). Baseline and 30-day ET-1 levels were similar between nesiritide and placebo groups ( $p > 0.05$ ).

**Conclusion:** Baseline ET-1 independently predicts 180-day mortality in ADHF. ET-1 may serve as a novel biomarker of mortality in ADHF independent of NT-proBNP.

